



FORM PTO-1449/And B (modified PTO/SB/08)				APPLICATION NO.: 10/582,028		ATTY. DOCKET NO.: H0498.70215US02	
<b>INFORMATION DISCLOSURE STATEMENT BY APPLICANT</b>				FILING DATE: June 8, 2006		CONFIRMATION NO.: Not Yet Assigned	
				APPLICANT: Howard Green et al.			
				GROUP ART UNIT: Not Yet Assigned		EXAMINER: Not Yet Assigned	
Sheet	1	of	4				

#### U.S. PATENT DOCUMENTS

Examiner's Initials #	Cite No.	U.S. Patent Document		Name of Patentee or Applicant of Cited Document	Date of Publication or Issue of Cited Document MM-DD-YYYY
		Number	Kind Code		

#### FOREIGN PATENT DOCUMENTS

Examiner's Initials #	Cite No.	Foreign Patent Document			Name of Patentee or Applicant of Cited Document	Date of Publication of Cited Document MM-DD-YYYY	Translation (Y/N)
		Office/ Country	Number	Kind Code			
	B1	WO	02/097068	A2	Inserm	12-05-2002	
	B2	WO	05/056765		President and Fellows of Harvard College	06-23-2005	

#### OTHER ART — NON PATENT LITERATURE DOCUMENTS

Examiner's Initials #	Cite No.	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	Translation (Y/N)
	C1	ALLEN-HOFFMANN, B.L. & RHEINWALD, J.G., Polycyclic aromatic hydrocarbon mutagenesis of human epidermal keratinocytes in culture. Proc Natl Acad Sci U S A. 1984 Dec;81(24):7802-6.	
	C2	BAGUTTI, C. et al., Dermal fibroblast-derived growth factors restore the ability of beta(1) integrin-deficient embryonal stem cells to differentiate into keratinocytes. Dev Biol. 2001 Mar 15;231(2):321-33.	
	C3	BAGUTTI, C. et al., Differentiation of embryonal stem cells into keratinocytes: comparison of wild-type and beta 1 integrin-deficient cells. Dev Biol. 1996 Oct 10;179(1):184-96.	
	C4	BANKS-SCHLEGEL, S. & GREEN, H.J. Involucrin synthesis and tissue assembly by keratinocytes in natural and cultured human epithelia. J Cell Biol. 1981 Sep;90(3):732-7.	
	C5	BARRANDON, Y. & GREEN, H., Three clonal types of keratinocyte with different capacities for multiplication. Proc Natl Acad Sci U S A. 1987 Apr;84(8):2302-6.	
	C6	CELLI, J. et al., Heterozygous germline mutations in the p53 homolog p63 are the cause of EEC syndrome. Cell. 1999 Oct 15;99(2):143-53.	
	C7	CORAUX, C. et al., Reconstituted skin from murine embryonic stem cells. Curr Biol. 2003 May 13;13(10):849-53.	
	C8	ECKERT, R. L. et al., The epidermis: genes on - genes off. J Invest Dermatol. 1997 Oct;109(4):501-9. Review.	
	C9	EVANS, M. J. & KAUFMAN, M. H., Establishment in culture of pluripotent cells from mouse embryos. Nature. 1981 Jul 9;292(5819):154-6.	
	C10	GALVIN, S. et al., Citizens Advice Bureaux in general practice: an illuminative evaluation. Health Soc Care Community. 2000 Jul;8(4):277-282.	

C11	ITSKOVITZ-ELDOR, J. et al., Differentiation of human embryonic stem cells into embryoid bodies compromising the three embryonic germ layers. <i>Mol Med.</i> 2000 Feb;6(2):88-95.	
C12	IUCHI, S. & GREEN, H., Nuclear localization of basonuclin in human keratinocytes and the role of phosphorylation. <i>Proc Natl Acad Sci U S A.</i> 1997 Jul 22;94(15):7948-53.	
C13	IUCHI, S. Three classes of C2H2 zinc finger proteins. <i>Cell Mol Life Sci.</i> 2001 Apr;58(4):625-35. Review.	
C14	IUCHI, S. et al., Alternative subcellular locations of keratinocyte basonuclin. <i>Exp Dermatol.</i> 2000 Jun;9(3):178-84.	
C15	KAUFMAN, D.S. et al., Hematopoietic colony-forming cells derived from human embryonic stem cells. <i>Proc Natl Acad Sci U S A.</i> 2001 Sep 11;98(19):10716-21. Epub 2001 Sep 4.	
C16	KEHAT, I. et al., Human embryonic stem cells can differentiate into myocytes with structural and functional properties of cardiomyocytes. <i>J Clin Invest.</i> 2001 Aug;108(3):407-14.	
C17	KUMAR, M. & MELTON, D., Pancreas specification: a budding question. <i>Curr Opin Genet Dev.</i> 2003 Aug;13(4):401-7. Review.	
C18	LI, W. et al., Signals that initiate, augment, and provide directionality for human keratinocyte motility. <i>J Invest Dermatol.</i> 2004 Oct;123(4):622-33.	
C19	LUMELSKY, N. et al., Differentiation of embryonic stem cells to insulin-secreting structures similar to pancreatic islets. <i>Science.</i> 2001 May 18;292(5520):1389-94. Epub 2001 Apr 26. Erratum in: <i>Science</i> 2001 Jul;293(5529):428.	
C20	MACINNES, M.A. et al., Differentiation and Enrichment of Functional Keratinocytes from Mouse Embryonic Stem (ES) Cells. Bethesda, MD, US, Vol. 11, no. SUPPL, 2000, page 410A, XP-001053766, ISSN: 1059-1524.	
C21	MAHONEY, M.G. et al., Translocation of the zinc finger protein basonuclin from the mouse germ cell nucleus to the midpiece of the spermatozoon during spermiogenesis. <i>Biol Reprod.</i> 1998 Aug;59(2):388-94.	
C22	MARTIN, G. R., Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. <i>Proc Natl Acad Sci U S A.</i> 1981 Dec;78(12):7634-8.	
C23	MCGRATH, J. A. et al., Hay-Wells syndrome is caused by heterozygous missense mutations in the SAM domain of p63. <i>Hum Mol Genet.</i> 2001 Feb 1;10(3):221-9.	
C24	MILLS, A. A. et al., p63 is a p53 homologue required for limb and epidermal morphogenesis. <i>Nature.</i> 1999 Apr 22;398(6729):708-13.	
C25	MOLL, R. et al., The catalog of human cytokeratins: patterns of expression in normal epithelia, tumors and cultured cells. <i>Cell.</i> 1982 Nov;31(1):11-24. Review.	
C26	O'CONNELL, J. T. et al., Identification of a basal/reserve cell immunophenotype in benign and neoplastic endometrium: a study with the p53 homologue p63. <i>Gynecol Oncol.</i> 2001 Jan;80(1):30-6.	
C27	PALMIERI, S. L. et al., Oct-4 transcription factor is differentially expressed in the mouse embryo during establishment of the first two extraembryonic cell lineages involved in implantation. <i>Dev Biol.</i> 1994 Nov;166(1):259-67.	
C28	PARENTEAU, N. L. et al., Induction of keratinocyte type-I transglutaminase in epithelial cells of the rat. <i>Differentiation.</i> 1986;33(2):130-41.	

C29	PARSA, R. et al., Association of p63 with proliferative potential in normal and neoplastic human keratinocytes. <i>J Invest Dermatol.</i> 1999 Dec;113(6):1099-105.	
C30	PELLEGRINI, G. et al., p63 identifies keratinocyte stem cells <i>Proc Natl Acad Sci U S A.</i> 2001 Mar 13;98(6):3156-61.	
C31	PHILLIPS, M. A. & RICE, R. H. Convergent differentiation in cultured rat cells from nonkeratinized epithelia: keratinocyte character and intrinsic differences. <i>J Cell Biol.</i> 1983 Sep;97(3):686-91.	
C32	QUINLAN, R. A. et al., Patterns of expression and organization of cytokeratin intermediate filaments. <i>Ann N Y Acad Sci.</i> 1985;455:282-306. Review.	
C33	REUBINOFF, B. E. et al., Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. <i>Nat Biotechnol.</i> 2000 Apr;18(4):399-404. Erratum in: <i>Nat Biotechnol</i> 2000 May;18(5):559.	
C34	RHEINWALD, J. G. & GREEN, H., Formation of a keratinizing epithelium in culture by a cloned cell line derived from a teratoma. <i>Cell.</i> 1975 Nov;6(3):317-30.	
C36	RHEINWALD, J. G. & GREEN, H., Serial cultivation of strains of human epidermal keratinocytes: the formation of keratinizing colonies from single cells. <i>Cell.</i> 1975 Nov;6(3):331-43.	
C36	RICE, R. H. & GREEN, H. Presence in human epidermal cells of a soluble protein precursor of the cross-linked envelope: activation of the cross-linking by calcium ions. <i>Cell.</i> 1979 Nov;18(3):681-94.	
C37	ROBERTSON, E. J. in <i>Teratocarcinomas and Embryonic Stem Cells: A Practical Approach</i> , Chapter 4: Embryo-derived stem cell lines, Oxford, UK: IRL Press, 1987; 71-112.	
C36	ROOP, D. R. et al., Transcriptional control of high molecular weight keratin gene expression in multistage mouse skin carcinogenesis. <i>Cancer Res.</i> 1988 Jun 1;48(11):3245-52.	
C34	ROSNER, M. H. et al., A POU-domain transcription factor in early stem cells and germ cells of the mammalian embryo. <i>Nature.</i> 1990 Jun 21;345(6277):686-92.	
C40	SCHOLER, H. R. et al., A family of octamer-specific proteins present during mouse embryogenesis: evidence for germline-specific expression of an Oct factor. <i>EMBO J.</i> 1989 Sep;8(9):2543-50.	
C41	SCHOLER, H. R. et al., Octamer binding proteins confer transcriptional activity in early mouse embryogenesis. <i>EMBO J.</i> 1989 Sep;8(9):2551-7.	
C42	SCHOLER, H. R. et al., Oct-4: a germline-specific transcription factor mapping to the mouse t-complex. <i>EMBO J.</i> 1990 Jul;9(7):2185-95.	
C43	SCHOLER, H. R. Octamania: the POU factors in murine development. <i>Trends Genet.</i> 1991 Oct;7(10):323-9. Review.	
C44	SCHULDINER, M. et al., Induced neuronal differentiation of human embryonic stem cells. <i>Brain Res.</i> 2001 Sep 21;913(2):201-5.	
C45	SCHULDINER, M. et al., Effects of eight growth factors on the differentiation of cells derived from human embryonic stem cells. <i>Proc Natl Acad Sci U S A.</i> 2000 Oct 10;97(21):11307-12.	
C46	SHAMBLOTT, M. J. et al., Human embryonic germ cell derivatives express a broad range of developmentally distinct markers and proliferate extensively in vitro. <i>Proc Natl Acad Sci U S A.</i> 2001 Jan 2;98(1):113-8.	

C47	SIGNORETTI, S. et al., p63 is a prostate basal cell marker and is required for prostate development. Am J Pathol. 2000 Dec;157(6):1769-75.	
C48	SIMON, M. & GREEN, H., Enzymatic cross-linking of involucrin and other proteins by keratinocyte particulates in vitro. Cell. 1985 Mar;40(3):677-83.	
C49	SMITH, A. G., Embryo-derived stem cells: of mice and men. Annu Rev Cell Dev Biol. 2001;17:435-62. Review.	
C50	THOMSON, J. A. & ODORICO, J. S., Human embryonic stem cell and embryonic germ cell lines. Trends Biotechnol. 2000 Feb;18(2):53-7.	
C51	THOMSON, J. A. et al., Isolation of a primate embryonic stem cell line. Proc Natl Acad Sci U S A. 1995 Aug 15;92(17):7844-8.	
C52	THOMSON, J. A. et al., Embryonic stem cell lines derived from human blastocysts. Science. 1998 Nov 6;282(5391):1145-7. Erratum in: Science 1998 Dec 4;282(5395):1827.	
C53	TIAN, Q. et al., Function of basonuclin in increasing transcription of the ribosomal RNA genes during mouse oogenesis. Development. 2001 Feb;128(3):407-16.	
C54	TOYOOKA, Y. et al., Embryonic stem cells can form germ cells in vitro. Proc Natl Acad Sci U S A. 2003 Sep 30;100(20):11457-62. Epub 2003 Sep 22.	
C55	TSENG, H. & GREEN, H., Association of basonuclin with ability of keratinocytes to multiply and with absence of terminal differentiation. J Cell Biol. 1994 Jul;126(2):495-506.	
C56	TSENG, H. & GREEN, H., Basonuclin: a keratinocyte protein with multiple paired zinc fingers. Proc Natl Acad Sci U S A. 1992 Nov 1;89(21):10311-5.	
C57	VAN BOKHOVEN, H. & MCKEON, F., Mutations in the p53 homolog p63: allele-specific developmental syndromes in humans. Trends Mol Med. 2002 Mar;8(3):133-9. Review.	
C58	VAN BOKHOVEN, H. et al., p63 Gene mutations in eec syndrome, limb-mammary syndrome, and isolated split hand-split foot malformation suggest a genotype-phenotype correlation. Am J Hum Genet. 2001 Sep;69(3):481-92. Epub 2001 Jul 17.	
C59	WATT, F. M. & GREEN, H., Stratification and terminal differentiation of cultured epidermal cells. Nature. 1982 Feb 4;295(5848):434-6.	
C60	WRIGHT, K. A. et al., Alternative delivery of keratinocytes using a polyurethane membrane and the implications for its use in the treatment of full-thickness burn injury. Burns. 1998 Feb;24(1):7-17.	
C61	YANG, A. et al., p63, a p53 homolog at 3q27-29, encodes multiple products with transactivating, death-inducing, and dominant-negative activities. Mol Cell. 1998 Sep;2(3):305-16.	
C62	YANG, A. et al., p63 is essential for regenerative proliferation in limb, craniofacial and epithelial development. Nature. 1999 Apr 22;398(6729):714-8.	

EXAMINER:

/Leon Lankford Jr/

DATE CONSIDERED:

03/29/2010

\* EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

\* a copy of this reference is not provided as it was previously cited by or submitted to the office in a prior application, Serial No. \_\_, filed \_\_, and relied upon for an earlier filing date under 35 U.S.C. 120 (continuation, continuation-in-part, and divisional applications).

[NOTE - No copies of U.S. patents, published U.S. patent applications, or pending, unpublished patent applications stored in the USPTO's Image File Wrapper (IFW) system, are included. See 37 CFR §1.98 and 1287OG163. Copies of all other patent(s), publication(s), unpublished, pending U.S. patent applications, or other information listed are provided as required by 37 CFR §1.98 unless 1) such copies were provided in an IDS in an earlier application that complies with 37 CFR §1.98, and 2) the earlier application is relied upon for an earlier filing date under 35 U.S.C. §120.]